

# The co-norm

## Unit 18

Gil Cohen

April 24, 2022

Throughout this unit, we let  $F/L$  be a function field extension of  $E/K$ .

## Definition 1

Let  $\mathfrak{p}$  be a prime divisor of  $E/K$ . To  $\mathfrak{p}$  we associate a divisor

$$\text{Con}(\mathfrak{p}) = \sum_{\mathfrak{P}/\mathfrak{p}} e(\mathfrak{P}/\mathfrak{p})\mathfrak{P}.$$

The map  $\text{Con}$  can be extended in the natural way to a group homomorphism

$$\text{Con} : \mathcal{D}(E/K) \rightarrow \mathcal{D}(F/L)$$

We call  $\text{Con}$  the **co-norm** and sometimes denote it by  $\text{Con}_{F/E}$ .

The following are straightforward properties of the co-norm.

## Lemma 2

- 1  $\alpha \geq 0 \implies \text{Con}(\alpha) \geq 0$ .
- 2 If  $\alpha, \beta$  are disjoint then so are  $\text{Con}(\alpha), \text{Con}(\beta)$ .
- 3 If  $F'/L'$  is an extension of  $F/L$  then

$$\text{Con}_{F'/E} = \text{Con}_{F'/F} \circ \text{Con}_{F/E}.$$

- 4  $\text{Con}$  is one to one.

The co-norm is a natural homomorphism in that it “lifts” a principle divisor of  $E$  to the corresponding divisor in  $F$ .

## Lemma 3

*For every  $x \in E^\times$ ,*

$$\begin{aligned}(x)_F &= \text{Con}((x)_E), \\(x)_{F,0} &= \text{Con}((x)_{E,0}), \\(x)_{F,\infty} &= \text{Con}((x)_{E,\infty}).\end{aligned}$$

Proof.

We start by proving that  $(x)_F = \text{Con}((x)_E)$ .

$$\begin{aligned}(x)_F &= \sum_{\mathfrak{P} \in \mathbb{P}(F/L)} v_{\mathfrak{P}}(x) \mathfrak{P} \\ &= \sum_{\mathfrak{p} \in \mathbb{P}(E/K)} \sum_{\mathfrak{P}/\mathfrak{p}} v_{\mathfrak{P}}(x) \mathfrak{P} \\ &= \sum_{\mathfrak{p} \in \mathbb{P}(E/K)} v_{\mathfrak{p}}(x) \sum_{\mathfrak{P}/\mathfrak{p}} e(\mathfrak{P}/\mathfrak{p}) \mathfrak{P} \\ &= \sum_{\mathfrak{p} \in \mathbb{P}(E/K)} v_{\mathfrak{p}}(x) \text{Con}(\mathfrak{p}) \\ &= \text{Con} \left( \sum_{\mathfrak{p} \in \mathbb{P}(E/K)} v_{\mathfrak{p}}(x) \mathfrak{p} \right) \\ &= \text{Con}((x)_E).\end{aligned}$$

Proof.

This proves the first item. Using it, we have that

$$(x)_F = \text{Con}((x)_E) = \text{Con}((x)_{E,0}) - \text{Con}((x)_{E,\infty}).$$

By the Lemma 2, the two divisors  $\text{Con}((x)_{E,0})$  and  $\text{Con}((x)_{E,\infty})$  are non-negative and disjoint, and the proof of items 2,3 follow. □

# Some notation

When we take a divisor  $\alpha \in \mathcal{D}(E/K)$  and would like to consider “it” over  $F/L$  we will identify  $\alpha$  with  $\text{Con}(\alpha)$ . To keep notation light, we will sometimes use  $\alpha$  rather than  $\text{Con}(\alpha)$ .

However, the degree and dimension of  $\alpha$  and  $\text{Con}(\alpha)$  typically will not be the same, and so we write  $\deg_F \alpha$  for  $\deg \text{Con}(\alpha)$  and  $\deg_E(\alpha)$  for  $\deg \alpha$ , and similarly define  $\dim_F \alpha$  as  $\dim \text{Con}(\alpha)$ .

# Degree under co-norm

## Theorem 4

*There exists a constant  $\lambda$  that depends only on  $F/E$  s.t.  $\forall \alpha \in \mathcal{D}(E/K)$ ,*

$$\deg_F \alpha = \lambda \cdot \deg_E \alpha.$$

*Moreover, if  $F/E$  is finite then  $\lambda = \frac{[F:E]}{[L:K]}$ .*

## Proof.

We first prove the theorem for the case that  $F/E$  is finite. It suffices to prove the theorem for prime divisors (as  $\deg_E$  and  $\deg_F = \deg \circ \text{Con}$  are homomorphisms).



# Degree under co-norm

Proof.

Take  $\mathfrak{p} \in \mathcal{D}(E/K)$ . Then,

$$\begin{aligned}\deg_F \mathfrak{p} &= \deg \operatorname{Con}(\mathfrak{p}) \\ &= \deg \sum_{\mathfrak{P}/\mathfrak{p}} e(\mathfrak{P}/\mathfrak{p}) \mathfrak{P} \\ &= \sum_{\mathfrak{P}/\mathfrak{p}} e(\mathfrak{P}/\mathfrak{p}) \deg \mathfrak{P}.\end{aligned}$$

Recall that

$$[L : K] \deg \mathfrak{P} = f(\mathfrak{P}/\mathfrak{p}) \deg_E \mathfrak{p},$$

and so, using the fundamental equality,

$$\deg_F \mathfrak{p} = \frac{\deg_E \mathfrak{p}}{[L : K]} \sum_{\mathfrak{P}/\mathfrak{p}} e(\mathfrak{P}/\mathfrak{p}) f(\mathfrak{P}/\mathfrak{p}) = \frac{[F : E]}{[L : K]} \cdot \deg_E \mathfrak{p}.$$

# Degree under co-norm

Proof.

We turn to consider the general case. It suffices to show that for all prime divisors  $p, q$ ,

$$\frac{\deg_E p}{\deg_F p} = \frac{\deg_E q}{\deg_F q}.$$

Assume towards a contradiction that for some prime divisors  $p, q$  it holds that

$$\frac{\deg_E p}{\deg_F p} < \frac{\deg_E q}{\deg_F q},$$

or, equivalently,

$$\frac{\deg_E p}{\deg_E q} < \frac{\deg_F p}{\deg_F q}.$$

# Degree under co-norm

Proof.

$$\frac{\deg_E p}{\deg_E q} < \frac{\deg_F p}{\deg_F q}.$$

Then, there are  $m, n \in \mathbb{N}$  s.t.

$$\frac{\deg_E p}{\deg_E q} < \frac{m}{n} < \frac{\deg_F p}{\deg_F q}$$

Equivalently,

$$\deg_F(np - mq) > 0,$$

$$\deg_E(np - mq) < 0.$$

# Degree under co-norm

Proof.

$$\deg_F(np - mq) > 0,$$

$$\deg_E(np - mq) < 0.$$

To get a contradiction, it suffices to prove that

$$\forall \mathfrak{a} \in \mathcal{D}(E/K) \quad \deg_E \mathfrak{a} > 0 \implies \deg_F \mathfrak{a} \geq 0.$$

Consider  $k \in \mathbb{N}$  sufficiently large (compared to  $g_E$ ). By Riemann-Roch,

$$\dim_E k\mathfrak{a} = k \deg_E \mathfrak{a} + 1 - g_E > 0.$$

Thus,  $\exists x \in E^\times$  s.t.  $(x)_E + k\mathfrak{a} \geq 0$ . By Lemma 2 and Lemma 3,

$$0 \leq \text{Con}((x)_E + k\mathfrak{a}) = (x)_F + k\text{Con}(\mathfrak{a}),$$

and so, as  $\deg((x)_F) = 0$ ,

$$\deg_F \mathfrak{a} = \deg \text{Con}(\mathfrak{a}) \geq 0.$$

